


Tedor Pharma
100ct. 50 mg/300 mg/40 mg
Butalbital, Acetaminophen,
and Caffeine Capsules, USP

Base Label



GRAPHICS PROOF

Rev.#	Date	BC Grade	Artist
1	01/08/16	A	KH
2	01/13/16	A	SS
3	01/28/16	A	JPC/CS

Size: 2.1250" x 5.0000"		Copy Position	THIS PROOF IS NOT INTENDED FOR COLOR REPRESENTATION. PLEASE REVIEW FOR COPY AND POSITIONING OF GRAPHICS AND TEXT.
Rev: 3			
			APPROVAL SIGNATURE/DATE: _____
FRONT BLACK	PMS 7461		RESEAL AREA
		#4	

Booklet

Page #2
(Back of Cover)

Page #3




Page #4

Page #5

Page #6

Back

Front

<p>Butalbital, Acetaminophen, and Caffeine Capsules, USP</p> <p>Rx only</p> <p>BOXED WARNING</p> <p>Hepatotoxicity</p> <p>Acetaminophen has been associated with cases of acute liver failure, at times resulting in liver transplant and death. Most of the cases of liver injury are associated with the use of acetaminophen at doses that exceed 4000 milligrams per day, and often involve more than one acetaminophen containing product.</p> <p>DESCRIPTION</p> <p>Butalbital, Acetaminophen, and Caffeine Capsules, USP are supplied in hard-gelatin capsule form for oral administration.</p> <p>Each capsule contains the following active ingredients:</p> <p>butalbital USP 50 mg acetaminophen USP 300 mg caffeine USP 40 mg</p> <p>Inactive Ingredients: croscollidone, microcrystalline cellulose, povidone, pregelatinized starch, stearic acid, talc, D&C yellow #10, FD&C yellow #6, FD&C green #3, FD&C red #40, gelatin, titanium dioxide, propylene glycol, hypromellose 2910 (6 MPAs), FD&C red #3, FD&C blue #1.</p>	<p>Butalbital (5-allyl-5-isobutylbarbituric acid), is a short to intermediate-acting barbiturate of molecular weight 224.26. It has the following structural formula:</p>  <p>C₁₁H₁₆N₂O₃ MW = 224.26</p> <p>Acetaminophen (4'-hydroxyacetanilide), is a non-opiate, non-salicylate analgesic and antipyretic of molecular weight 151.16. It has the following structural formula:</p>  <p>C₉H₉NO₂ MW = 151.16</p>	<p>Caffeine (1,3,7-trimethylxanthine), is a central nervous system stimulant of molecular weight 194.19. It has the following structural formula:</p>  <p>C₈H₁₀N₄O₂ MW = 194.19</p> <p>CLINICAL PHARMACOLOGY</p> <p>This combination drug product is intended as a treatment for tension headache.</p> <p>It consists of a fixed combination of butalbital, acetaminophen, and caffeine. The role each component plays in the relief of the complex of symptoms known as tension headache is incompletely understood.</p> <p>Pharmacokinetics</p> <p>The behavior of the individual components is described below.</p>	<p>Butalbital</p> <p>Butalbital is well absorbed from the gastrointestinal tract and is expected to distribute to most tissues in the body. Barbiturates in general may appear in breast milk and readily cross the placental barrier. They are bound to plasma and tissue proteins to a varying degree and binding increases directly as a function of lipid solubility.</p> <p>Elimination of butalbital is primarily via the kidney (59% to 88% of the dose) as unchanged drug or metabolites. The plasma half-life is about 35 hours. Urinary excretion products include parent drug (about 3.6% of the dose), 5-isobutyl-5-(2,3-dihydroxypropyl) barbituric acid (about 24% of the dose), 5-allyl-5-(3-hydroxy-2-methyl-1-propyl) barbituric acid (about 4.8% of the dose), products with the barbituric acid ring hydrolyzed with excretion of urea (about 14% of the dose), as well as unidentified materials. Of the material excreted in the urine, 32% is conjugated.</p> <p>The <i>in vitro</i> plasma protein binding of butalbital is 45% over the concentration range of 0.5-20 mg/mL. This falls within the range of plasma protein binding (20%-45%) reported with other barbiturates such as phenobarbital, pentobarbital, and secobarbital sodium. The plasma-to-blood concentration ratio was almost unity, indicating that there is no preferential distribution of butalbital into either plasma or blood cells.</p> <p>See OVERDOSAGE for toxicity information.</p>	<p>Acetaminophen</p> <p>Acetaminophen is rapidly absorbed from the gastrointestinal tract and is distributed throughout most body tissues. The plasma half-life is 1.25 to 3 hours, but may be increased by liver damage and following overdose. Elimination of acetaminophen is principally by liver metabolism (conjugation) and subsequent renal excretion of metabolites. Approximately 85% of an oral dose appears in the urine within 24 hours of administration, most as the glucuronide conjugate, with small amounts of other conjugates and unchanged drug.</p> <p>See OVERDOSAGE for toxicity information.</p> <p>Caffeine</p> <p>Like most xanthines, caffeine is rapidly absorbed and distributed in all body tissues and fluids, including the CNS, fetal tissues, and breast milk.</p> <p>Caffeine is cleared through metabolism and excretion in the urine. The plasma half-life is about 3 hours. Hepatic biotransformation prior to excretion results in about equal amounts of 1-methylxanthine and 1-methyluric acid. Of the 70% of the dose that is recovered in the urine, only 3% is unchanged drug.</p> <p>See OVERDOSAGE for toxicity information.</p>	<p>INDICATIONS AND USAGE</p> <p>Butalbital, acetaminophen, and caffeine capsules are indicated for the relief of the symptom complex of tension (or muscle contraction) headache.</p> <p>Evidence supporting the efficacy and safety of this combination product in the treatment of multiple recurrent headaches is unavailable. Caution in this regard is required because butalbital is habit-forming and potentially abusable.</p> <p>CONTRAINDICATIONS</p> <p>This product is contraindicated under the following conditions:</p> <ul style="list-style-type: none"> Hypersensitivity or intolerance to any component of this product Patients with porphyria <p>WARNINGS</p> <p>Butalbital is habit-forming and potentially abusable. Consequently, the extended use of this product is not recommended.</p> <p>Hepatotoxicity</p> <p>Acetaminophen has been associated with cases of acute liver failure, at times resulting in liver transplant and death. Most of the cases of liver injury are associated with the use of acetaminophen at doses that exceed 4000 milligrams per day, and often involve more than one acetaminophen containing product. The excessive intake of acetaminophen may be intentional</p>	<p>to cause self-harm or unintentional as patients attempt to obtain more pain relief or unknowingly take other acetaminophen-containing products.</p> <p>The risk of acute liver failure is higher in individuals with underlying liver disease and in individuals who ingest alcohol while taking acetaminophen. Instruct patients to look for acetaminophen or APAP on package labels and not to use more than one product that contains acetaminophen. Instruct patients to seek medical attention immediately upon ingestion of more than 4000 milligrams of acetaminophen per day, even if they feel well.</p> <p>Serious skin reactions</p> <p>Rarely, acetaminophen may cause serious skin reactions such as acute generalized exanthematous pustulosis (AGEP), Stevens-Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. Patients should be informed about the signs of serious skin reactions, and use of the drug should be discontinued at the first appearance of skin rash or any other sign of hypersensitivity.</p> <p>Hypersensitivity / anaphylaxis</p> <p>There have been post-marketing reports of hypersensitivity and anaphylaxis associated with use of acetaminophen. Clinical signs included swelling of the face, mouth, and throat, respiratory distress, urticaria, rash, pruritus, and vomiting. There were infrequent reports of life-threatening anaphylaxis</p>	<p>requiring emergency medical attention. Instruct patients to discontinue Butalbital, Acetaminophen, and Caffeine Capsules, USP immediately and seek medical care if they experience these symptoms. Do not prescribe Butalbital, Acetaminophen, and Caffeine Capsules, USP for patients with acetaminophen allergy.</p> <p>PRECAUTIONS</p> <p>General</p> <p>Butalbital, acetaminophen, and caffeine capsules should be prescribed with caution in certain special-risk patients, such as the elderly or debilitated, and those with severe impairment of renal or hepatic function, or acute abdominal conditions.</p> <p>Information for Patients</p> <p>This product may impair mental and/or physical abilities required for the performance of potentially hazardous tasks such as driving a car or operating machinery. Such tests should be avoided while taking this product.</p> <p>Alcohol and other CNS depressants may produce an additive CNS depression when taken with this combination product, and should be avoided.</p> <p>Butalbital may be habit-forming. Patients should take the drug only for as long as it is prescribed, in the amounts prescribed, and no more frequently than prescribed.</p>	<p>For information on use in geriatric patients, see PRECAUTIONS/Geriatric Use.</p> <ul style="list-style-type: none"> Do not take Butalbital, Acetaminophen, and Caffeine Capsules, USP if you are allergic to any of its ingredients. If you develop signs of allergy such as a rash or difficulty breathing stop taking Butalbital, Acetaminophen, and Caffeine Capsules, USP and contact your healthcare provider immediately. Do not take more than 4000 milligrams of acetaminophen per day. Call your doctor if you took more than the recommended dose. <p>Laboratory Tests</p> <p>In patients with severe hepatic or renal disease, effects of therapy should be monitored with serial liver and/or renal function tests.</p> <p>Drug Interactions</p> <p>The CNS effects of butalbital may be enhanced by monoamine oxidase (MAO) inhibitors.</p> <p>Butalbital, acetaminophen, and caffeine may enhance the effects of: other narcotic analgesics, alcohol, general anesthetics, tranquilizers such as chloriazepoxide, sedative-hypnotics, or other CNS depressants, causing increased CNS depression.</p>	<p>Drug/Laboratory Test Interactions</p> <p>Acetaminophen may produce false-positive test results for urinary 5-hydroxyindoleacetic acid.</p> <p>Carcinogenesis, Mutagenesis, Impairment of Fertility</p> <p>No adequate studies have been conducted in animals to determine whether acetaminophen or butalbital have a potential for carcinogenesis, mutagenesis or impairment of fertility.</p> <p>Pregnancy</p> <p>Teratogenic Effects</p> <p><i>Pregnancy Category C:</i> Animal reproduction studies have not been conducted with this combination product. It is also not known whether butalbital, acetaminophen, and caffeine can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. This product should be given to a pregnant woman only when clearly needed.</p> <p>Nonteratogenic Effects</p> <p>Withdrawal seizures were reported in a two-day-old male infant whose mother had taken a butalbital-containing drug during the last two months of pregnancy. Butalbital was found in the infant's serum. The infant was given phenobarbital 5 mg/kg, which was tapered without further seizure or other withdrawal symptoms.</p>
---	--	--	---	---	--	---	--	--	--

19.9063"

<p>Nursing Mothers</p> <p>Caffeine, barbiturates, and acetaminophen are excreted in breast milk in small amounts, but the significance of their effects on nursing infants is not known. Because of potential for serious adverse reactions in nursing infants from butalbital, acetaminophen, and caffeine, a decision should be made whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.</p> <p>Pediatric Use</p> <p>Safety and effectiveness in pediatric patients below the age of 12 have not been established.</p> <p>Geriatric Use</p> <p>Clinical studies of butalbital, acetaminophen and caffeine capsules did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.</p> <p>Butalbital is known to be substantially excreted by the kidney, and the risk</p>	<p>of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.</p> <p>ADVERSE REACTIONS</p> <p>Frequently Observed</p> <p>The most frequently reported adverse reactions are drowsiness, lightheadedness, dizziness, sedation, shortness of breath, nausea, vomiting, abdominal pain, and intoxicated feeling.</p> <p>Infrequently Observed</p> <p>All adverse events tabulated below are classified as infrequent.</p> <p>Central Nervous System: headache, shaky feeling, tingling, agitation, fainting, fatigue, heavy eyelids, high energy, hot spells, numbness, sluggishness, seizure. Mental confusion, excitement, or depression can also occur due to intolerance, particularly in elderly or debilitated patients, or due to overdose of butalbital.</p> <p>Autonomic Nervous System: dry mouth, hyperhidrosis.</p> <p>Gastrointestinal: difficulty swallowing, heartburn, flatulence, constipation.</p>	<p>Cardiovascular: tachycardia.</p> <p>Musculoskeletal: leg pain, muscle fatigue.</p> <p>Genitourinary: diuresis.</p> <p>Miscellaneous: pruritus, fever, earache, nasal congestion, tinnitus, euphoria, allergic reactions.</p> <p>Several cases of dermatological reactions, including toxic epidermal necrolysis and erythema multiforme, have been reported.</p> <p>The following adverse drug events may be borne in mind as potential effects of the components of this product. Potential effects of high dosage are listed in the OVERDOSAGE section.</p> <p>Acetaminophen: allergic reactions, rash, thrombocytopenia, agranulocytosis.</p> <p>Caffeine: cardiac stimulation, irritability, tremor, dependence, nephrotoxicity, hyperglycemia.</p> <p>DRUG ABUSE AND DEPENDENCE</p> <p>Abuse and Dependence</p> <p>Butalbital</p> <p><i>Barbiturates may be habit-forming.</i> Tolerance, psychological dependence, and physical dependence may occur especially following prolonged use of</p>	<p>high doses of barbiturates. The average daily dose for the barbiturate addict is usually about 1500 mg. As tolerance to barbiturates develops, the amount needed to maintain the same level of intoxication increases; tolerance to a fatal dosage, however, does not increase more than two-fold. As this occurs, the margin between an intoxication dosage and fatal dosage becomes smaller. The lethal dose of a barbiturate is far less if alcohol is also ingested. Major withdrawal symptoms (convulsions and delirium) may occur within 16 hours and last up to 5 days after abrupt cessation of these drugs. Intensity of withdrawal symptoms gradually declines over a period of approximately 15 days. Treatment of barbiturate dependence consists of cautious and gradual withdrawal of the drug. Barbiturate-dependent patients can be withdrawn by using a number of different withdrawal regimens. One method involves initiating treatment at the patient's regular dosage level and gradually decreasing the daily dosage as tolerated by the patient.</p> <p>OVERDOSAGE</p> <p>Following an acute overdose of butalbital, acetaminophen, and caffeine, toxicity may result from the barbiturate or the acetaminophen. Toxicity due to caffeine is less likely, due to the relatively small amounts in this formulation.</p> <p>Signs and Symptoms</p> <p>Toxicity from <i>barbiturate</i> poisoning includes drowsiness, confusion, and coma; respiratory depression; hypotension; and hypovolemic shock.</p>	<p>potential risk of hepatotoxicity; acetaminophen levels drawn less than 4 hours post-ingestion may be misleading. To obtain the best possible outcome, NAC should be administered as soon as possible where impending or evolving liver injury is suspected. Intravenous NAC may be administered when circumstances preclude oral administration.</p> <p>DOSE AND ADMINISTRATION</p> <p>One or 2 capsules every 4 hours as needed. Total daily dosage should not exceed 6 capsules.</p> <p>Extended and repeated use of this product is not recommended because of the potential for physical dependence.</p> <p>HOW SUPPLIED</p> <p>Butalbital, Acetaminophen, and Caffeine Capsules USP, 50 mg/300 mg/40 mg</p> <p>Containing 50 mg butalbital, 300 mg acetaminophen, and 40 mg caffeine. Available as hard gelatin capsules with a green cap, printed with "Tedor/255", and a yellow body, printed with "Tedor/255". The capsules are supplied in bottles of 100 capsules (NDC 42195-955-10). Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Dispense in a tight, light-resistant container.</p>	<p>Rx only</p> <p>Manufactured for: Xspire Pharma Ridgeland, MS 39157</p> <p>Manufactured by: Tedor Pharma Inc. Cumberland, RI 02864 Rev. 01/16</p>	<p>NDC 42195-955-10</p> <p>Butalbital, Acetaminophen, and Caffeine Capsules, USP 50 mg/300 mg/40 mg</p> <p>⚠️ ONLY 100 CAPSULES</p> <p>DESCRIPTION</p> <p>Each capsule contains: Butalbital USP 50 mg Acetaminophen USP 300 mg Caffeine USP 40 mg</p> <p>USUAL ADULT DOSAGE: 1 or 2 capsules every four hours. Total daily dose should not exceed 6 capsules. See package insert for additional prescribing information.</p> <p>DISPENSE in a tight, light-resistant container as defined in the USP.</p> <p>STORE at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].</p> <p>KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.</p> <p>Manufactured for: Xspire Pharma, LLC Ridgeland, MS 39157 www.xspirex.com 13001 Rev. 01/16</p>  <p>Lift Here</p>
--	--	---	--	---	--	---

Page #7

Page #8
GRAPHICS PROOF

Page #9

Page #10

Cover
(Page #1)

Size: 2.1250" x 5.0000"						<p>THIS PROOF IS NOT INTENDED FOR COLOR REPRESENTATION. PLEASE REVIEW FOR COPY AND POSITIONING OF GRAPHICS AND TEXT.</p> <p>PLATINUM PRESS FAX: (469)733-1510</p> <p>APPROVAL SIGNATURE/DATE: _____</p> <p>APPROVAL SIGNATURE/DATE: _____</p>
Rev: 3						
						<p>#4</p>
FRONT BLACK	PMS 7461	BACK BLACK				

APPROVED REVISE AND RE-PROOF